

IVIG NEUROLOGY REFERRAL FORM

3070 McCann Farm Drive | Suite 101 Garnet Valley, PA 19060

Toc	lay'	's [Da:	е

1-866-317-0672 TEL: 610-545-6040 FAX: 610-545-6030

	CURRENT PATIENT
1	NEW PATIENT

Medical History: Cardiac Diseculergies Current Medications (if necessal PRACTICE NAME	y, please fax a comple	te list)	IACT INFORMATION		RMATION
	PEFFI	RRAL SOURCE INFORM	MATION		
##				#	
#		##		#	
###		##		# #	
nsured's Name ligible for Medicare 🏻 Yes 🗀	No If yes, Medicare#	Relation to Pa Presc	cription Card \square Yes \square N	No If Yes, Carrier	
el n#	. Fax Pcn#	Policy/Gro RXID#	oup# RX	Group#	
CD-10 Diagnosis Code G61.81 Chronic Inflammatory G61.9 Inflammatory Polyneuro G70.01 Myasthenia Gravis wi M33.20 Polymyositis, organ inv	Demyelinating Polyneu opathy, unspecified th (Acute) Exacerbation	uropathy (CIDP)	G70.80 Lambert-Eaton S M36.0 Dermatomyositis G25.82 Stiff-Person Syndi G35 Multiple Sclerosis (<i>N</i> ICD-10: DX:	rome (S)	ed
Is this the first dose? Yes					
	FUSION PUMP: days, as a loading d grams	ose, then gro			
List product	days, as a loading d gramsadryl) 25-50 mg orally	ose, then gro			
List product	days, as a loading days, as a lo	before infusion before infusion pefore infusion N SPECIALTY CARE Pl kg: mg orally, IM or IV a phenhydramine 50m pp infusion, call 911,	ROTOCOL nd decrease the rate ag, orally, IM or IV and give Epinephrine 0.3n	of infusion. contact physicia	_ cycle(s)
List product	cadryl) 25-50 mg orally by than 66 pounds or 30 hiphenhydramine 50 ristop infusion, give Digreathing problem: stog IV or IM. Begin NSS necessary for children	before infusion before infusion N SPECIALTY CARE Plants May be a composite of the second se	ROTOCOL nd decrease the rate ng, orally, IM or IV and give Epinephrine 0.3n f 100-150ml/hr and co	of infusion. contact physicia ng/0.3ml subcutar ntact physician.	n neously,
List product	days, as a loading layer. EEACTIONS: PER ELWY than 66 pounds or 30 piphenhydramine 50 restop infusion, give Digreathing problem: stop IV or IM. Begin NSS arecessary for children layer. If Epinephrine is a days of the days are layer.	before infusion before infusion before infusion before infusion before infusion before infusion before infusion. N SPECIALTY CARE Plants are supply to the plants of the	ROTOCOL nd decrease the rate ng, orally, IM or IV and give Epinephrine 0.3n f 100-150ml/hr and co 6 pounds: Diphenhyd 5ml 1:1000 subcutane	of infusion. contact physicia ng/0.3ml subcutar ntact physician.	n neously,